

United States Patent [19]

Rheinheimer et al.

[11] Patent Number: 4,983,210

[45] Date of Patent: Jan. 8, 1991

[54] **ISOXAZOLINES, THEIR PREPARATION AND THEIR USE**

[75] Inventors: **Joachim Rheinheimer**, Ludwigshafen; **Karl Eicken**, Wachenheim; **Hans Theobald**, Limburgerhof; **Thomas Kuekenhoehner**, Frankenthal; **Karl-Otto Westphalen**, Speyer; **Bruno Wuerzer**, Otterstadt; **Juergen Frank**, Schwetzingen; **Norbert Meyer**, Ladenburg, all of Fed. Rep. of Germany

[73] Assignee: **BASF Aktiengesellschaft**, Ludwigshafen, Fed. Rep. of Germany

[21] Appl. No.: 326,986

[22] Filed: Mar. 22, 1989

[30] Foreign Application Priority Data

Mar. 23, 1988 [DE] Fed. Rep. of Germany 3809765

[51] Int. Cl.⁵ C07D 261/04; C07D 403/04; C07D 403/12; A01N 43/80

[52] U.S. Cl. 71/94; 71/88; 546/209; 546/275; 548/240

[58] Field of Search 71/88, 94; 546/275, 546/209; 548/240

[56] References Cited

U.S. PATENT DOCUMENTS

3,631,169 12/1971 Minami et al. 544/137

FOREIGN PATENT DOCUMENTS

0187345 7/1986 European Pat. Off. .

2724677 12/1978 Fed. Rep. of Germany .

OTHER PUBLICATIONS

Cozzi et al., *Gazzetta Chimica Italiana*, vol. 116, pp. 717-719 (1986).

Houk et al., *J. Am. Chem. Soc.*, vol. 106, pp. 3880-3882 (1984).

Annunziata et al., *Tetrahedron*, vol. 42, No. 7, pp. 2129-2134 (1986).


Primary Examiner—Mukund J. Shah

Assistant Examiner—E. Bernhardt

Attorney, Agent, or Firm—Keil and Weinkauff

[57] **ABSTRACT**

Isoxazolines of the general formula I



where the substituents have the following meanings:

R¹ is C₁-C₁₂-alkyl, C₃-C₇-cycloalkyl, substituted or unsubstituted phenyl, or a saturated or unsaturated 5- or 6-membered heterocycle;

R², R³, R⁴, R⁵ and R⁶ are hydrogen, C₁-C₄-alkyl or benzyl and


R⁷ is substituted or unsubstituted C₂-C₆-alkenyl, C₅-C₇-cycloalkenyl, phenyl, naphthyl or thienyl,

processes for their manufacture, and their use.

3 Claims, No Drawings

ISOXAZOLINES, THEIR PREPARATION AND THEIR USE

The present invention relates to isoxazolines of the general formula I



where R¹ is C₁-C₁₂-alkyl, C₃-C₇-cycloalkyl, phenyl which can carry one to five halogen atoms and/or one to three of the groups C₁-C₄-alkyl, C₁-C₄-alkoxy and/or C₁-C₄-haloalkyl, or a saturated or unsaturated 5-membered or 6-membered heterocyclic structure containing a nitrogen, oxygen or sulfur atom as a ring member, R², R³, R⁴, R⁵ and R⁶ are each hydrogen, C₁-C₄-alkyl or benzyl and R⁷ is C₂-C₆-alkenyl or C₅-C₇-cycloalkenyl, where these groups may carry one to three phenyl radicals and/or halogen atoms, or is phenyl which may carry one to three of the substituents C₁-C₄-alkyl, C₁-C₄-alkoxy, C₁-C₄-haloalkyl, phenyl or phenoxy, cyano and/or halogen, or is naphthyl or thienyl, where these aromatic rings may carry one to three C₁-C₄-alkyl groups and/or halogen atoms.


The present invention furthermore relates to processes for the preparation of the compounds I, herbicides which contain these compounds and methods for controlling undesirable plant growth with the compounds I.

DE-A 27 24 677 describes herbicidal 2-(benzyloxy-methyl)-tetrahydrofurans. However, because of the poor selectivity with respect to weeds and the relatively high application rates, their action is unsatisfactory.

It is an object of the present invention to provide compounds having improved herbicidal properties.

We have found that this object is achieved by the isoxazolines I defined at the outset. We have furthermore found processes for the preparation of the compounds and methods for controlling undesirable plant growth with these novel compounds.

The compounds I are obtained, for example, by etherifying an appropriately substituted 5-hydroxymethylisoxazoline derivative II with a compound III in a conventional manner in an inert organic solvent in the presence of a base to give an isoxazoline I.



In formula III, Nu is a nucleophilic leaving group, such as chlorine, bromine, iodine, arylsulfonyl or alkyl-



sulfonyl, e.g. toluenesulfonyl, or another equivalent leaving group. Compounds of the formula III having a reactive substituent Nu are known from the literature or are readily obtainable on the basis of the general technical knowledge.

This reaction is generally carried out at from 0° to 120° C., preferably from 10° to 80° C., in an aprotic solvent, preferably in an ether, such as diethyl ether, methyl tert-butyl ether, tetrahydrofuran or dioxane, or dimethyl sulfoxide, dimethylformamide or a mixture of these.

Examples of suitable bases are alkali metal or alkaline earth metal hydrides, such as NaH or CaH₂, alkali metal hydroxides, such as NaOH or KOH, alkali metal carbonates, such as Na₂CO₃ or K₂CO₃, alkali metal amides, such as sodium amide or lithium diisopropylamide, and tertiary amines, such as triethylamine, diisopropylethylamine, N,N-dimethyl-p-aminopyridine, N-methylpyrrolidine, N,N,N',N'-tetramethylethylenediamine, 1,5-diazabicyclo[4.3.0]non-5-ene and 1,8-diazabicyclo[5.4.0]undec-7-ene. The reaction is preferably carried out in the presence of one of the abovementioned inorganic bases.

The reaction can be accelerated by adding a phase-transfer catalyst, such as a crown ether or a quaternary ammonium salt, if heat-sensitive radicals of the educts necessitate mild reaction conditions.

The 5-hydroxymethylisoxazoline derivatives II required for the reaction are obtained in a conventional manner (C. Grundmann, Synthesis 1970, 347 and J. M. J. Trochet, S. Jaccard-Thorndahl, L. Faivre and R. Massard, Helv. Chim. Acta 56 (1973), 1303) from the corresponding oximes IV by reaction with an alkyl alcohol of the formula V,



With regard to the use of the novel isoxazolines I in accordance with regulations, preferred substituents are the following radicals: R¹ is alkyl, such as methyl, isopropyl, n-butyl, isobutyl, tert-butyl, pentyl, 1-methylbutyl, 2-methylbutyl, 3-methylbutyl, 1,1-dimethylpropyl, 1,2-dimethylpropyl, 2,2-dimethylpropyl, 1-ethylpropyl, hexyl, 1-methylpentyl, 2-methylpentyl, 3-methylpentyl, 4-methylpentyl, 1,1-dimethylbutyl, 1,2-dimethylbutyl, 1,3-dimethylbutyl, 2,2-dimethylbutyl, 2,3-dimethylbutyl, 3,3-dimethylbutyl, 1-ethylbutyl, 2-ethylbutyl, 1,1,2-trimethylpropyl, 1,2,2-trimethylpropyl, 1-ethyl-1-methylpropyl and 1-ethyl-2-methylpropyl, in particular methyl, ethyl, propyl, 1-methylethyl, butyl, 1-methylpropyl, 2-methylpropyl and 1,1-dimethylethyl; cycloalkyl, such as cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl and cycloheptyl, in particular cyclopropyl, cyclopentyl and cyclohexyl; phenyl which may be sub-

stituted by one to five, in particular one to three, halogen atoms, such as fluorine, chlorine, bromine and/or iodine, in particular fluorine, chlorine or bromine, and/or by one to three of the abovementioned C₁-C₄-alkyl groups, in particular methyl and ethyl, alkoxy groups, such as methoxy, ethoxy, propoxy, 1-methylethoxy, butoxy, 1-methylpropoxy, 2-methylpropoxy or 1,1-dimethylethoxy, in particular methoxy or ethoxy, and/or haloalkyl groups, such as fluoromethyl, difluoromethyl, trifluoromethyl, chlorodifluoromethyl, dichlorofluoromethyl, trichloromethyl, 1-fluoroethyl, 2-fluoroethyl, 2,2-difluoroethyl, 2,2,2-trifluoroethyl, 2-chloro-2,2-difluoroethyl, 2,2-dichloro-2-fluoroethyl, 2,2,2-trichloroethyl or pentafluoroethyl, in particular difluoromethyl or trifluoromethyl; a saturated or unsaturated 5-membered or 6-membered heterocyclic structure, such as tetrahydrofuran, tetrahydrothiophene, pyrrolidine, tetrahydropyran, tetrahydrothiopyran, piperidine, furan, thienyl, pyrrole or pyridine, in particular tetrahydropyran, furfuryl, thienyl, pyridyl or tetrahydrofuran; R² is benzyl or C₁-C₄-alkyl as stated under R¹, in particular hydrogen, methyl or ethyl; R³ is in general and in particular one of the radicals stated under R²; R⁴ is in general one of the radicals stated under R², in particular hydrogen, methyl, ethyl, propyl, 1-methylethyl or benzyl; R⁵ and R⁶ are each in general and in particular one of the radicals stated under R²; and R⁷ is alkenyl, such as ethenyl, 1-propenyl, 2-propenyl, 1-methylethenyl, 1-butenyl, 2-butenyl, 3-butenyl, 1-methyl-1-propenyl, 1-methyl-2-propenyl, 2-methyl-1-propenyl, 2-methyl-2-propenyl, 1-pentenyl, 2-pentenyl, 3-pentenyl, 4-pentenyl, 1-methyl-1-butenyl, 2-methyl-1-butenyl, 3-methyl-1-butenyl, 1-methyl-2-butenyl, 2-methyl-2-butenyl, 3-methyl-2-butenyl, 1-methyl-3-butenyl, 2-methyl-3-butenyl, 3-methyl-3-butenyl, 1,1-dimethyl-2-propenyl, 1,1-dimethyl-1-propenyl, 1,1-dimethyl-2-propenyl, 1,2-dimethyl-1-propenyl, 1,2-dimethyl-2-propenyl, 1-ethyl-1-propenyl, 1-ethyl-2-propenyl, 1-hexenyl, 2-hexenyl, 3-hexenyl, 4-hexenyl, 5-hexenyl, 1-methyl-1-pentenyl, 2-methyl-1-pentenyl, 3-methyl-1-pentenyl, 4-methyl-1-pentenyl, 1-methyl-2-pentenyl, 2-methyl-2-pentenyl, 3-methyl-2-pentenyl, 4-methyl-2-pentenyl, 1-methyl-3-pentenyl, 2-methyl-3-pentenyl, 3-methyl-3-pentenyl, 4-methyl-3-pentenyl, 1-methyl-4-pentenyl, 2-methyl-4-pentenyl, 3-methyl-4-pentenyl, 4-methyl-4-pentenyl, 1,1-dimethyl-2-butenyl, 1,1-dimethyl-3-butenyl, 1,2-dimethyl-1-butenyl, 1,2-dimethyl-2-butenyl, 1,2-dimethyl-3-butenyl, 1,3-dimethyl-1-butenyl, 1,3-dimethyl-2-butenyl, 1,3-dimethyl-3-butenyl, 2,2-dimethyl-3-butenyl, 2,3-dimethyl-1-butenyl, 2,3-dimethyl-2-butenyl, 2,3-dimethyl-3-butenyl, 3,3-dimethyl-1-butenyl, 1-ethyl-1-butenyl, 1-ethyl-2-butenyl, 1-ethyl-3-butenyl, 2-ethyl-1-butenyl, 2-ethyl-2-butenyl, 2-ethyl-3-butenyl, 1,2,2-trimethyl-2-propenyl, 1-ethyl-1-methyl-2-propenyl, 1-ethyl-2-methyl-1-propenyl or 1-ethyl-2-methyl-2-propenyl, in particular ethenyl, 1-methylethylenyl, 2-methyl-1-propenyl or 2-methyl-1-butenyl; C₅-C₇-cycloalkenyl, such as cyclopentenyl, cyclohexenyl or cycloheptenyl, in particular cyclopentenyl or cyclohexenyl; or phenyl, naphthyl or thienyl, and when R⁷ is alkenyl or cycloalkenyl it can be substituted by, in particular, phenyl and/or one to three of the halogen atoms stated under R¹, when R⁷ is phenyl it can be substituted by one to three of the abovementioned halogen atoms, C₁-C₄-haloalkyl groups and/or a phenyl radical, a phenoxy radical and/or cyano groups, and when R⁷ is naphthyl or thienyl it can be substituted by

one to three of the abovementioned C₁-C₄-alkyl groups and/or halogen atoms.

Particularly preferred compounds I are those in which R¹ is straight-chain or branched C₁-C₅-alkyl, C₃-C₆-cycloalkyl, tetrahydropyran, pyridyl, furyl, phenyl, chlorophenyl, methoxyphenyl or methylphenyl, R² and R³ are each hydrogen, methyl or ethyl, R⁴ is hydrogen, straight-chain or branched C₁-C₄-alkyl or benzyl, R⁵ and R⁶ are each hydrogen or methyl and R⁷ is phenyl, mono-, di- or trichlorophenyl, mono-, di- or trifluorophenyl, mono-, di- or trimethylphenyl, fluorochlorophenyl, 2-fluoro-6-methylphenyl, 2,4-difluoro-6-methylphenyl, 2-chloro-6-methylphenyl, bromophenyl, mono- or diethylphenyl, phenoxyphenyl, methoxyphenyl, cyanophenyl, trifluoromethylphenyl, naphthyl, thienyl, mono- or dichlorothiophenyl, methylthienyl, 1-methylethylenyl, 1- and 2-methylpropenyl, chloroethenyl or chloropropenyl.

The isoxazolines I, or herbicidal agents containing them, may be applied for instance in the form of directly sprayable solutions, powders, suspensions (including high-percentage aqueous, oily or other suspensions), dispersions, emulsions, oil dispersions, pastes, dusts, broadcasting agents, or granules by spraying, atomizing, dusting, broadcasting or watering. The forms of application depend entirely on the purpose for which the agents are being used, but they must ensure as fine a distribution of the active ingredients according to the invention as possible.

For the preparation of solutions, emulsions, pastes and oil dispersions to be sprayed direct, mineral oil fractions of medium to high boiling point, such as kerosene or diesel oil, further coal-tar oils, and oils of vegetable or animal origin, aliphatic, cyclic and aromatic hydrocarbons such as benzene, toluene, xylene, paraffin, tetrahydronaphthalene, alkylated naphthalenes and their derivatives such as methanol, ethanol, propanol, butanol, chloroform, carbon tetrachloride, cyclohexanol, cyclohexanone, chlorobenzene, isophorone, etc., and strongly polar solvents such as dimethylformamide, dimethyl sulfoxide, N-methylpyrrolidone, water, etc. are suitable.

Aqueous formulations may be prepared from emulsion concentrates, pastes, oil dispersions or wettable powders by adding water. To prepare emulsions, pastes and oil dispersions the ingredients as such or dissolved in an oil or solvent may be homogenized in water by means of wetting or dispersing agents, adherents or emulsifiers. Concentrates which are suitable for dilution with water may be prepared from active ingredient, wetting agent, adherent, emulsifying or dispersing agent and possibly solvent or oil.

Examples of surfactants are: alkali metal, alkaline earth metal and ammonium salts of ligninsulfonic acid, naphthalenesulfonic acids, phenolsulfonic acids, alkylaryl sulfonates, alkyl sulfates, and alkyl sulfonates, alkali metal and alkaline earth metal salts of dibutyl naphthalenesulfonic acid, lauryl ether sulfate, fatty alcohol sulfates, alkali metal and alkaline earth metal salts of fatty acids, salts of sulfated hexadecanols, heptadecanols, and octadecanols, salts of sulfated fatty alcohol glycol ethers, condensation products of sulfonated naphthalene and naphthalene derivatives with formaldehyde, condensation products of naphthalene or naphthalenesulfonic acids with phenol and formaldehyde, polyoxyethylene octylphenol ethers, ethoxylated isooctylphenol ethoxylated octylphenol and ethoxylated nonylphenol, alkylphenol polyglycol ethers, tributylphenyl

polyglycol ethers, alkylaryl polyether alcohols, iso-tridecyl alcohol, fatty alcohol ethylene oxide condensates, ethoxylated castor oil, polyoxyethylene alkyl ethers, ethoxylated polyoxypropylene, lauryl alcohol polyglycol ether acetal, sorbitol esters, lignin, sulfite waste liquors and methyl cellulose.

Powders, dusts and broadcasting agents may be prepared by mixing or grinding the active ingredients with a solid carrier.

Granules, e.g., coated, impregnated or homogeneous granules, may be prepared by bonding the active ingredients to solid carriers. Examples of solid carriers are mineral earths such as silicic acid, silica gels, silicates, talc, kaolin, attapulugus clay, limestone, lime, chalk, bole, loess, clay, dolomite, diatomaceous earth, calcium sulfate, magnesium sulfate, magnesium oxide, ground plastics, fertilizers such as ammonium sulfate, ammonium phosphate, ammonium nitrate, and ureas, and vegetable products such as grain flours, bark meal, wood meal, and nutshell meal, cellulosic powders, etc.

The formulations contain from 0.1 to 95, and preferably 0.5 to 90, % by weight of active ingredients.

Examples of formulations are given below.

I. 90 parts by weight of compound no. 1 is mixed with 10 parts by weight of N-methyl-alpha-pyrrolidone. A mixture is obtained which is suitable for application in the form of very fine drops.

II. 20 parts by weight of compound no. 10 is dissolved in a mixture consisting of 80 parts by weight of xylene, 10 parts by weight of the adduct of 8 to 10 moles of ethylene oxide and 1 mole of oleic acid-N-monoethanolamide, 5 parts by weight of the calcium salt of dodecylbenzenesulfonic acid, and 5 parts by weight of the adduct of 40 moles of ethylene oxide and 1 mole of castor oil. By pouring the solution into 100,000 parts by weight of water and uniformly distributing it therein, an aqueous dispersion is obtained containing 0.02% by weight of the active ingredient.

III. 20 parts by weight of compound no. 17 is dissolved in a mixture consisting of 40 parts by weight of cyclohexanone, 30 parts by weight of isobutanol, 20 parts by weight of the adduct of 7 moles of ethylene oxide and 1 mole of isooctylphenol, and 10 parts by weight of the adduct of 40 moles of ethylene oxide and 1 mole of castor oil. By pouring the solution into 100,000 parts by weight of water and finely distributing it therein, an aqueous dispersion is obtained containing 0.2% by weight of the active ingredient.

IV. 20 parts by weight of compound no. 29 is dissolved in a mixture consisting of 25 parts by weight of cyclohexanol, 65 parts by weight of a mineral oil fraction having a boiling point between 210° and 280° C., and 10 parts by weight of the adduct of 40 moles of ethylene oxide and 1 mole of castor oil. By pouring the solution into 100,000 parts by weight of water and uniformly distributing it therein, an aqueous dispersion is obtained containing 0.02% by weight of the active ingredient.

V. 20 parts by weight of compound no. 50 is well mixed with 3 parts by weight of the sodium salt of diisobutyl-naphthalene-alpha-sulfonic acid, 17 parts by weight of the sodium salt of a lignin-sulfonic acid obtained from a sulfite waste liquor, and 60 parts by weight of powdered silica gel, and triturated in a hammer mill. By uniformly distributing the mixture in 20,000 parts by weight of water, a spray liquor is obtained containing 0.1% by weight of the active ingredient.

VI. 3 parts by weight of compound no. 72 is intimately mixed with 97 parts by weight of particulate kaolin. A dust is obtained containing 3% by weight of the active ingredient.

VII. 30 parts by weight of compound no. 121 is intimately mixed with a mixture consisting of 92 parts by weight of powdered silica gel and 8 parts by weight of paraffin oil which has been sprayed onto the surface of this silica gel. A formulation of the active ingredient is obtained having good adherence.

VIII. 20 parts of compound no. 125 is intimately mixed with 2 parts of the calcium salts of dodecylbenzenesulfonic acid, 8 parts of a fatty alcohol polyglycol ether, 2 parts of the sodium salt of a phenolsulfonic acid-urea-formaldehyde condensate and 68 parts of a paraffinic mineral oil. A stable oily dispersion is obtained.

The active ingredients may be applied pre- or post-emergence. If certain crop plants tolerate the active ingredients less well, application techniques may be used in which the herbicidal agents are sprayed from suitable equipment in such a manner that the leaves of sensitive crop plants are if possible not touched, and the agents reach the soil or the unwanted plants growing beneath the crop plants (post-directed, lay-by treatment).

The application rates depend on the objective to be achieved, the time of the year, the plants to be combated and their growth stage, and are from 0.001 to 3.0, preferably 0.01 to 1.0, kg of active ingredient per hectare.

In view of the number of application methods possible, the compounds according to the invention, or agents containing them, may be used in a further large number of crops for removing unwanted plants. The following crops are given by way of example:

Botanical name	Common name
<i>Allium cepa</i>	onions
<i>Ananas comosus</i>	pineapples
<i>Arachis hypogaea</i>	peanuts (groundnuts)
<i>Asparagus officinalis</i>	asparagus
<i>Avena sativa</i>	oats
<i>Beta vulgaris</i> spp. <i>altissima</i>	sugarbeets
<i>Beta vulgaris</i> spp. <i>rapa</i>	fodder beets
<i>Beta vulgaris</i> spp. <i>esculenta</i>	table beets, red beets
<i>Brassica napus</i> var. <i>napus</i>	rapeseed
<i>Brassica napus</i> var. <i>napobrassica</i>	swedes
<i>Brassica napus</i> var. <i>rapa</i>	turnips
<i>Brassica rapa</i> var. <i>silvestris</i>	
<i>Camellia sinensis</i>	tea plants
<i>Carthamus tinctorius</i>	safflower
<i>Carya illinoensis</i>	pecan trees
<i>Citrus limon</i>	lemons
<i>Citrus maxima</i>	grapefruits
<i>Citrus reticulata</i>	mandarins
<i>Citrus sinensis</i>	orange trees
<i>Coffea arabica</i> (<i>Coffea canephora</i> , <i>Coffea liberica</i>)	coffee plants
<i>Cucumis melo</i>	melons
<i>Cucumis sativus</i>	cucumbers
<i>Cynodon dactylon</i>	Bermudagrass
<i>Daucus carota</i>	carrots
<i>Elais guineensis</i>	oil palms
<i>Fragaria vesca</i>	strawberries
<i>Glycine max</i>	soybeans
<i>Gossypium hirsutum</i> (<i>Gossypium arboreum</i> , <i>Gossypium herbaceum</i> , <i>Gossypium vitifolium</i>)	cotton
<i>Helianthus annuus</i>	sunflowers
<i>Helianthus tuberosus</i>	Jerusalem arti- choke
<i>Hevea brasiliensis</i>	rubber plants

-continued

Botanical name	Common name
<i>Hordeum vulgare</i>	barley
<i>Humulus lupulus</i>	hops
<i>Ipomoea batatas</i>	sweet potatoes
<i>Juglans regia</i>	walnut trees
<i>Lactuca sativa</i>	lettuce
<i>Lens culinaris</i>	lentils
<i>Linum usitatissimum</i>	flax
<i>Lycopersicon lycopersicum</i>	tomatoes
<i>Malus</i> spp.	apple trees
<i>Manihot esculenta</i>	cassava
<i>Medicago sativa</i>	alfalfa (lucerne)
<i>Mentha piperita</i>	peppermint
<i>Musa</i> spp.	banana plants
<i>Nicotiana tabacum</i> (<i>N. rustica</i>)	tobacco
<i>Olea europaea</i>	olive trees
<i>Oryza sativa</i>	rice
<i>Panicum miliaceum</i>	millet
<i>Phaseolus lunatus</i>	limabeans
<i>Phaseolus mungo</i>	mungbeans
<i>Phaseolus vulgaris</i>	snapbeans, green beans, dry beans
<i>Pennisetum glaucum</i>	pearl millet
<i>Petroselinum crispum</i> spp. <i>tuberosum</i>	parsley
<i>Picea abies</i>	Norway spruce
<i>Abies alba</i>	fir trees
<i>Pinus</i> spp.	pine trees
<i>Pisum sativum</i>	English peas
<i>Prunus avium</i>	cherry trees
<i>Prunus domestica</i>	plum trees
<i>Prunus dulcis</i>	almond trees
<i>Prunus persica</i>	peach trees
<i>Pyrus communis</i>	pear trees
<i>Ribes sylvestre</i>	redcurrants
<i>Ribes uva-crispa</i>	gooseberries
<i>Ricinus communis</i>	castor-oil plants
<i>Saccharum officinarum</i>	sugar cane
<i>Secale cereale</i>	rye
<i>Sesamum indicum</i>	sesame
<i>Solanum tuberosum</i>	Irish potatoes
<i>Sorghum bicolor</i> (<i>s. vulgare</i>)	sorghum
<i>Sorghum dochna</i>	sorgo
<i>Spinacia oleracea</i>	spinach
<i>Theobroma cacao</i>	cacao plants
<i>Trifolium pratense</i>	red clover
<i>Triticum aestivum</i>	wheat
<i>Triticum durum</i>	durum wheat
<i>Vaccinium corymbosum</i>	blueberries
<i>Vaccinium vitis-idaea</i>	cranberries
<i>Vicia faba</i>	tick beans
<i>Vigna sinensis</i> (<i>V. unguiculata</i>)	cow peas
<i>Vitis vinifera</i>	grapes
<i>Zea mays</i>	Indian corn, sweet corn, maize

To increase the spectrum of action and to achieve synergistic effects, the isoxazolines of the formula I may be mixed and applied together with numerous representatives of other herbicidal or growth-regulating active ingredients groups. Examples of suitable components are diazines, 4H-3,1-benzoxazine derivative, benzothiadiazinones, 2,6-dinitroanilines, N-phenylcarbamates, thiolcarbamates, halocarboxylic acids, triazines, amides, ureas, diphenyl ethers, triazinones, uracils, benzofuran derivatives, cyclohexane-1,3-dione derivatives,

quinolinecarboxylic acid derivatives, phenoxy- or heteroaryloxy-phenylpropionic acids and salts, esters and amides thereof, etc.

It may also be useful to apply the novel compounds of the formula I, either alone or in combination with other herbicides, in admixture with other crop protection agents, e.g., agents for combating pests or phytopathogenic fungi or bacteria. The compounds may also be mixed with solutions of mineral salts used to remedy nutritional or trace element deficiencies. Non-phytotoxic oils and oil concentrates may also be added.

SYNTHESIS EXAMPLES

The directions given in the synthesis examples below were employed, after appropriate modifications to the starting materials, to obtain further compounds of the formula I; the compounds obtained are listed in the tables below with physical data. Those compounds for which no data are given may be produced analogously from the appropriate materials. In view of their close structural similarity with the compounds produced and investigated, they are expected to have a similar action.

EXAMPLE 1

Manufacturing example for a starting material of the formula II: 5-hydroxymethyl-5-methyl-3-phenyl-4,5-dihydro-1,2-oxazole


At 10° C., 380 g of a 10% strength sodium hypochloride solution in which 0.3 g of sodium hydroxide is additionally dissolved is added dropwise to 50.0 g of benzaldoxime and 37.2 g of 2-methyl-2-propenol in 80 ml of diethyl ether. The mixture is stirred overnight at room temperature, the organic phase is separated off and extracted 5 times, each time with 100 ml of diethyl ether, and the extracts are dried over sodium sulfate and evaporated down. There remains 72.3 g of crude product, which is distilled at 140° C./0.01 mbar. Melting point: 49°-50° C. Further 4,5-dihydro-1,2-oxazoles may be prepared analogously.

EXAMPLE 2

General manufacturing example for isoxazolines of the formula I:


0.12 mol of sodium hydride is added to 50 ml of a 1:1 mixture of tetrahydrofuran and dimethylformamide. At room temperature, a solution of 0.10 ml of a 5-hydroxymethyl-4,5-dihydro-1,2-oxazole in 80 ml of a 1:1 mixture of tetrahydrofuran and dimethylformamide is then dripped in and the mixture is stirred for 1 hour at 60° C. Subsequently, a solution of 0.1 mol of the particular benzyl or allyl halide in 20 ml of the same solvent mixture is dripped in at room temperature. In many cases, the mixture has to be stirred for a further 2 hours at 60° C. to complete the conversion. The mixture is then stirred into 1 liter of ice water, extracted, dried and evaporated down. The crude product may be purified further by chromatography over silica gel or distillation under reduced pressure.

TABLE



No.	R ¹	R ²	R ³	R ⁴	R ⁵	R ⁶	R ⁷	Phys. data
1	methyl	H	H	CH ₃	H	H	phenyl	
2	methyl	H	H	CH ₃	H	H	2-methylphenyl	
3	methyl	H	H	CH ₃	H	H	2-fluorophenyl	
4	ethyl	H	H	CH ₃	H	H	phenyl	
5	ethyl	H	H	CH ₃	H	H	2-methylphenyl	¹ H: 1.13(t), 1.38(s), 2.20-2.40(m), 2.35(s)
6	ethyl	H	H	CH ₃	H	H	2-chlorophenyl	
7	ethyl	H	H	CH ₃	H	H	2-chloro-6-fluorophenyl	<i>n</i> _D ²⁴ = 1.5102
8	ethyl	H	H	CH ₃	H	H	2-fluorophenyl	
9	isopropyl	H	H	CH ₃	H	H	phenyl	
10	isopropyl	H	H	CH ₃	H	H	2-chlorophenyl	
11	isopropyl	H	H	CH ₃	H	H	2-fluorophenyl	¹ H: 1.10(d), 1.37(s), 2.60(d), 3.00(d), 4.55(s) <i>n</i> _D ²³ = 1.5058 <i>n</i> _D ²³ = 1.5425 <i>n</i> _D ²³ = 1.5214 <i>n</i> _D ²⁴ = 1.5106 <i>n</i> _D ²³ = 1.4731 <i>n</i> _D ²⁴ = 1.4854 <i>n</i> _D ²³ = 1.5594 <i>n</i> _D ²³ = 1.5035 <i>n</i> _D ²² = 1.5145
12	isopropyl	H	H	CH ₃	H	H	2-chloro-6-fluorophenyl	
13	isopropyl	H	H	CH ₃	H	H	3-phenoxyphenyl	
14	isopropyl	H	H	CH ₃	H	H	2-cyanophenyl	
15	isopropyl	H	H	CH ₃	H	H	2-methoxyphenyl	
16	isopropyl	H	H	CH ₃	H	H	2-trifluoromethylphenyl	
17	isopropyl	H	H	CH ₃	H	H	2,4-difluorophenyl	
18	isopropyl	H	H	CH ₃	H	H	1-naphthyl	
19	isopropyl	H	H	CH ₃	H	H	2,4,6-trimethylphenyl	
20	isopropyl	H	H	CH ₃	H	H	2-thienyl	
21	isopropyl	H	H	CH ₃	H	H	3-chloro-2-thienyl	
22	isopropyl	H	H	CH ₃	H	H	2-chloro-4-thienyl	
23	isopropyl	H	H	CH ₃	H	H	1-methylethenyl	<i>n</i> _D ²⁴ = 1.4554
24	isopropyl	H	H	CH ₃	H	H	2-methylpropen-1-yl	<i>n</i> _D ²³ = 1.4604
25	isopropyl	H	H	CH ₃	H	H	2-chloropropen-1-yl	
26	1-methylpropyl	H	H	CH ₃	H	H	phenyl	
27	1-methylpropyl	H	H	CH ₃	H	H	2-chlorophenyl	
28	1-methylpropyl	H	H	CH ₃	H	H	2-fluorophenyl	
29	1-methylpropyl	H	H	CH ₃	H	H	2-methylphenyl	¹ H: 0.80-1.00(m), 1.43(s), 3.40-3.55(m), 4.57(cm). <i>n</i> _D ²⁴ = 1.5030
30	1-methylpropyl	H	H	CH ₃	H	H	2-chloro-6-fluorophenyl	
31	1-methylpropyl	H	H	CH ₃	H	H	2,4-difluorophenyl	
32	2-methylpropyl	H	H	CH ₃	H	H	phenyl	
33	2-methylpropyl	H	H	CH ₃	H	H	2-methylphenyl	<i>n</i> _D ²⁴ = 1.5023
34	2-methylpropyl	H	H	CH ₃	H	H	2-chlorophenyl	
35	2-methylpropyl	H	H	CH ₃	H	H	2-fluorophenyl	
36	2-methylpropyl	H	H	CH ₃	H	H	2-chloro-6-fluorophenyl	<i>n</i> _D ²⁴ = 1.5031
37	cyclopropyl	H	H	CH ₃	H	H	phenyl	
38	cyclopropyl	H	H	CH ₃	H	H	2-methylphenyl	<i>n</i> _D ²³ = 1.9256
39	cyclopropyl	H	H	CH ₃	H	H	2-chlorophenyl	
40	cyclopropyl	H	H	CH ₃	H	H	2-fluorophenyl	
41	cyclopropyl	H	H	CH ₃	H	H	2,4-difluorophenyl	
42	cyclopropyl	H	H	CH ₃	H	H	2-chloro-6-fluorophenyl	<i>n</i> _D ²³ = 1.5231
43	cyclopentyl	H	H	CH ₃	H	H	phenyl	
44	cyclopentyl	H	H	CH ₃	H	H	2-methylphenyl	<i>n</i> _D ²⁴ = 1.5232
45	cyclopentyl	H	H	CH ₃	H	H	2-chlorophenyl	
46	cyclopentyl	H	H	CH ₃	H	H	2-fluorophenyl	
47	cyclopentyl	H	H	CH ₃	H	H	2-chloro-6-fluorophenyl	<i>n</i> _D ²⁴ = 1.5210
48	cyclopentyl	H	H	CH ₃	H	H	2,4-difluorophenyl	
49	phenyl	H	H	CH ₃	H	H	phenyl	
50	phenyl	H	H	CH ₃	H	H	2-methylphenyl	<i>n</i> _D ²⁴ = 1.5710
51	phenyl	H	H	CH ₃	H	H	2-chlorophenyl	
52	phenyl	H	H	CH ₃	H	H	2-chloro-6-fluorophenyl	¹ H: 1.47(s), 2.95(d), 3.45(d) 4.75 (s)
53	phenyl	H	H	CH ₃	H	H	2-fluorophenyl	
54	4-tetrahydropyran-2-yl	H	H	CH ₃	H	H	phenyl	
55	4-tetrahydropyran-2-yl	H	H	CH ₃	H	H	2-methylphenyl	<i>n</i> _D ²³ = 1.5246
56	4-tetrahydropyran-2-yl	H	H	CH ₃	H	H	2-chlorophenyl	
57	4-tetrahydropyran-2-yl	H	H	CH ₃	H	H	2-fluorophenyl	
58	4-tetrahydropyran-2-yl	H	H	CH ₃	H	H	2-chloro-6-fluorophenyl	<i>n</i> _D ²³ = 1.5236 <i>n</i> _D ²³ = 1.5234
59	3-tetrahydropyran-2-yl	H	H	CH ₃	H	H	2-methylphenyl	
60	methyl	H	H	H	CH ₃	H	2-chloro-6-fluorophenyl	bp.0.4 = 158-160
61	methyl	H	H	H	CH ₃	H	2,6-dichlorophenyl	bp.0.2 = 158-160
62	methyl	H	H	H	CH ₃	H	2-chlorophenyl	bp.0.3 = 150-152
63	ethyl	H	H	H	H	H	4-chloro-2-fluorophenyl	bp.0.4 = 163-165
64	ethyl	H	H	H	H	H	2,3,6-trichlorophenyl	bp.0.3 = 208-210
65	ethyl	H	H	H	H	H	2-chlorophenyl	bp.0.3 = 158-160
66	ethyl	H	H	H	H	H	2,6-dichlorophenyl	bp.0.2 = 183-185
67	ethyl	H	H	H	H	H	2-chloro-6-fluorophenyl	bp.0.2 = 157-158
68	ethyl	H	H	H	H	H	phenyl	bp.0.4 = 148-150
69	ethyl	H	H	H	H	H	2-methylphenyl	bp.0.4 = 162-164

TABLE-continued



No.	R ¹	R ²	R ³	R ⁴	R ⁵	R ⁶	R ⁷	Phys. data
70	methyl	H	H	H	H	H	phenyl	bp.0.1 = 126-128
71	methyl	H	H	H	H	H	2-methylphenyl	bp.0.06 = 146-148
72	methyl	H	H	H	H	H	2-chloro-6-fluorophenyl	bp.0.05 = 166-168
73	methyl	H	H	H	H	H	2,6-dichlorophenyl	bp.2 = 192-194
74	methyl	H	H	H	H	H	2-chlorophenyl	bp.0.05 = 156-158
75	methyl	H	H	H	H	H	4-chloro-2-fluorophenyl	bp.0.3 = 156-158
76	methyl	H	H	H	H	H	2,3,6-trichlorophenyl	bp.0.5 = 208-210
77	methyl	H	H	H	H	H	2-chloro-4-fluorophenyl	bp.0.06 = 160-162
78	methyl	H	H	H	H	H	4-fluorophenyl	bp.0.05 = 144-146
79	methyl	H	H	H	H	H	2,4-dichlorophenyl	bp.0.4 = 172-174
80	methyl	H	H	H	H	H	3,4-dichlorophenyl	bp.1.2 = 184-187
81	isopropyl	H	H	CH ₃	H	H	2,6-difluorophenyl	n _D ²³ = 1.4848
82	isopropyl	H	H	CH ₃	H	H	2,6-dimethylphenyl	n _D ²³ = 1.5075
83	1-methylpropyl	H	H	CH ₃	H	H	3-chloro-2-thienyl	n _D ²⁴ = 1.5045
84	1-methylpropyl	H	H	CH ₃	H	H	2-chloro-4-thienyl	n _D ²⁴ = 1.5240
85	1-methylpropyl	H	H	CH ₃	H	H	2-trifluoromethylphenyl	n _D ²⁴ = 1.4716
86	3-tetrahydropyranyl	H	H	CH ₃	H	H	2-chloro-6-fluorophenyl	n _D ²³ = 1.5212
87	3-tetrahydropyranyl	H	H	CH ₃	H	H	2-fluorophenyl	
88	n-propyl	H	H	CH ₃	H	H	phenyl	
89	n-propyl	H	H	CH ₃	H	H	2-methylphenyl	
90	n-propyl	H	H	CH ₃	H	H	2-chloro-6-fluorophenyl	
91	n-propyl	H	H	CH ₃	H	H	2-fluorophenyl	
92	n-propyl	H	H	CH ₃	H	H	2-bromophenyl	
93	t-butyl	H	H	CH ₃	H	H	2-chloro-6-fluorophenyl	n _D ²² = 1.5010
94	t-butyl	H	H	CH ₃	H	H	2-methylphenyl	n _D ²² = 1.5020
95	t-butyl	H	H	CH ₃	H	H	2-fluorophenyl	
96	t-butyl	H	H	CH ₃	H	H	2-ethylphenyl	
97	t-butyl	H	CH ₃	CH ₃	H	H	2-methylphenyl	
98	phenyl	CH ₃	CH ₃	CH ₃	H	H	2-methylphenyl	
99	isopropyl	H	H	ethyl	H	H	2-methylphenyl	n _D ²⁴ = 1.5050
100	isopropyl	H	H	ethyl	H	H	2-chloro-6-fluorophenyl	n _D ²⁴ = 1.5036
101	isopropyl	H	H	ethyl	H	CH ₃	2-chloro-6-fluorophenyl	
102	isopropyl	H	H	benzyl	H	H	2-methylphenyl	
103	isopropyl	H	H	benzyl	H	H	2-chloro-6-fluorophenyl	n _D ²⁴ = 1.5420
104	isopropyl	H	H	benzyl	H	H	2-fluorophenyl	
105	isopropyl	H	H	i-propyl	H	H	2-methylphenyl	n _D ²⁴ = 1.5049
106	isopropyl	H	H	i-propyl	H	H	2-chloro-6-fluorophenyl	n _D ²⁴ = 1.5035
107	2-furyl	H	H	CH ₃	H	H	2-chloro-6-fluorophenyl	
108	2-furyl	H	H	CH ₃	H	H	2-methylphenyl	
109	2-thienyl	H	H	CH ₃	H	H	2-methylphenyl	
110	3-pyridyl	H	H	CH ₃	H	H	2-methylphenyl	
111	3-pyridyl	H	H	CH ₃	H	H	2-chloro-6-fluorophenyl	
112	2-chlorophenyl	H	H	CH ₃	H	H	2-methylphenyl	
113	2-chlorophenyl	H	H	CH ₃	H	H	2-chloro-6-fluorophenyl	n _D ²⁴ = 1.5705
114	3-chlorophenyl	H	H	CH ₃	H	H	2-methylphenyl	
115	4-chlorophenyl	H	H	CH ₃	H	H	2-fluorophenyl	
116	2-methoxyphenyl	H	H	CH ₃	H	H	2-methylphenyl	
117	2-methoxyphenyl	H	H	CH ₃	H	H	2-chloro-6-fluorophenyl	
118	2-methoxyphenyl	H	H	CH ₃	H	H	phenyl	
119	isopropyl	H	H	n-propyl	H	H	2-methylphenyl	
120	n-butyl	H	H	CH ₃	H	H	2-chloro-6-fluorophenyl	
121	isopropyl	H	H	CH ₃	H	H	2-methylphenyl	¹ H: 1.10(d), 1.37(s), 2.30(s), 4.60(mc)
122	isopropyl	H	H	H	H	H	4-fluorophenyl	¹ H: 1.15(d), 2.60-3.10(m), 4.55(s)
123	isopropyl	H	H	H	H	H	2-methylphenyl	¹ H: 1.12(d), 2.32(s), 2.60-3.10(m) 3.45-3.65(m), 4.55(s) 4.60-4.80(m).
124	methyl	H	H	H	H	H	2-fluorophenyl	n _D ²⁴ = 1.5094
125	isopropyl	H	H	CH ₃	H	CH ₃	phenyl	n _D ²³ = 1.4850
126	2-chlorophenyl	H	H	CH ₃	H	H	2-fluorophenyl	n _D ²⁵ = 1.5642
127	2-methoxyphenyl	H	H	CH ₃	H	H	2-chlorophenyl	n _D ²³ = 1.5784
128	isopropyl	H	H	benzyl	H	H	phenyl	n _D ²⁴ = 1.5425

*refractive indexes (n_D), boiling points (Kp) in C.° or selected ¹H-NMR signals: δ in ppm using tetramethylsilane as internal standard (¹H).

USE EXAMPLES

The action of the isoxazolines of the formula I on plant growth is demonstrated by the following greenhouse experiments:

The vessels employed were plastic flowerpots having a volume of 300 cm³ and filled with a sandy loam containing about 3.0% humus. The seeds of the test plants were sown separately, according to species.

For the preemergence treatment, the active ingredients were applied to the surface of the soil immediately after the seeds had been sown. The compounds were emulsified or suspended in water as vehicle, and sprayed through finely distributing nozzles. After the agents had been applied, the vessels were lightly sprinkler-irrigated to induce germination and growth. Transparent plastic covers were then placed on the vessels until the plants had taken root. The cover ensured uniform germination of the plants, insofar as this was not impaired by the active ingredients. The application rates were 0.5 and 1.0 kg/ha.

For the postemergence treatment, the plants were grown, depending on growth form, to a height of 3 to 15 cm before being treated. In this treatment method, either plants which had been sown in the pots and grown there were selected, or they were cultivated separately as seedlings and transplanted to the pots a few days before being treated. The application rate for postemergence treatment was 1.0 kg/ha. No covers were placed on the vessels in this method.

The pots were set up in the greenhouse, species from warmer climates in warmer areas (20° to 35° C.) and species from moderate climates at 10° to 20° C. The experiments were run for from 2 to 4 weeks. During this time the plants were tended and their reactions to the various treatments assessed. The assessment scale was 0 to 100, 100 denoting nonemergence or complete destruction of at least the visible plant parts, and 0 denoting no damage or normal growth.

The plants used in the greenhouse experiments were:

Abbrev.	Botanical name	Common name
BRNSW	<i>Brassica napus</i>	rapeseed
BROIN	<i>Bromus inermis</i>	smooth broome
CHEAL	<i>Chenopodium album</i>	lambsquarters (goosefoot)
ELEIN	<i>Eleusine indica</i>	
ECHCG	<i>Echinochloa crus-galli</i>	barnyardgrass
HELAN	<i>Helianthus annuus</i>	sunflowers
LOLMU	<i>Lolium multiflorum</i>	annual ryegrass
POAAN	<i>Poa annua</i>	annual bluegrass


The compounds of Examples 7 and 12, applied pre-emergence at a rate of 1.0 kg/ha, combated unwanted plants extremely well, and were well tolerated by rape.

Compound no. 52, applied postemergence at a rate of 1.0 kg/ha, had a herbicidal action on *Chenopodium album* without damaging crop plants.

Compound no. 42, applied preemergence at a rate of 0.5 kg/ha, had a very good herbicidal action of broad-leaved plants without damaging, for instance, sunflowers.

We claim:

1. An isoxazoline of the formula I




where the substituents have the following meanings:

R¹ is phenyl substituted by from one to five halogens and from one to three of the groups C₁-C₄-alkyl, C₁-C₄-alkoxy or C₁-C₄-haloalkyl; or phenyl substituted by from one to five halogens or by from one to three of the groups C₁-C₄-alkyl, C₁-C₄-alkoxy or C₁-C₄-haloalkyl

R², R³, R⁴, R⁵ and R⁶ are hydrogen, C₁-C₄-alkyl or benzyl and

R⁷ is C₂-C₆-alkenyl or C₅-C₇-cycloalkenyl, or C₂-C₆-alkenyl or C₅-C₇-cycloalkenyl substituted by one to three phenyl radicals and halogen atoms, or C₂-C₆-alkenyl or C₅-C₇-cycloalkenyl substituted by one to three; phenyl radicals or halogen atoms; unsubstituted phenyl or phenyl substituted by from one to three of the following substituents: C₁-C₄-alkyl, C₁-C₄-alkoxy, C₁-C₄-haloalkyl, phenyl, phenoxy, cyano and halogen; or unsubstituted naphthyl or thienyl, or naphthyl or thienyl substituted by from one to three C₁-C₄-alkyl groups and halogen atoms, or naphthyl or thienyl substituted by from one to three C₁-C₄-alkyl groups or halogen atoms.

2. A process for combating the growth of unwanted plants, wherein the unwanted plants or their habitat are treated with a herbicidally effective amount of an isoxazoline of the formula I



where the substituents have the following meanings:

R¹ is C₁-C₁₂-alkyl, C₃-C₇-cycloalkyl, unsubstituted phenyl or phenyl substituted by from one to five halogens and from one to three of the groups C₁-C₄-alkyl, C₁-C₄-alkoxy or C₁-C₄-haloalkyl; or phenyl substituted by from one to five halogens or by from one to three of the group C₁-C₄-alkyl, C₁-C₄-alkoxy or C₁-C₄-haloalkyl; or R¹ is tetrahydrofuranyl, tetrahydrothiophenyl, pyrrolidinyl, tetrahydropyranyl, tetrahydrothiopyranyl, piperidyl, furanyl, thienyl, pyrrolyl or pyridyl directly attached to the 3-position of the isoxazoline ring;

R², R³, R⁴, R⁵ and R⁶ are hydrogen, C₁-C₄-alkyl or benzyl and

R⁷ is C₂-C₆-alkenyl or C₅-C₇-cycloalkenyl, or C₂-C₆-alkenyl or C₅-C₇-cycloalkenyl substituted by one to three phenyl radicals or halogen atoms; unsubstituted phenyl or phenyl substituted by from one to three of the following substituents: C₁-C₄-alkyl, C₁-C₄-alkoxy, C₁-C₄-haloalkyl, phenyl, phenoxy, cyano and halogen; or unsubstituted naphthyl or thienyl, or naphthyl or thienyl substituted by from one to three C₁-C₄-alkyl groups and halogen atoms, or naphthyl or thienyl substituted by one to three C₁-C₄-alkyl groups or halogen atoms.

3. A herbicidal composition containing a herbicidally effective amount of the isoxazoline of the formula I in claim 2, along with an inert carrier or diluent.

* * * * *